

University in St. Louis, School of Social Work, Saint Louis, MO/US, ³New York University School of Medicine, Department of Psychiatry, New York, NY/US

Background: The 2010 global burden of disease study indicated that mental and behavioral disorders account for approximately 9.5% of all disability-adjusted life years and 23.6% of all years lived with disability in China. With rapid socioeconomic transformation, the Chinese mental health system is in great need of service expansion with delivery models capable of operationalizing treatment packages into local practices. For Chinese cities facing many unique challenges today, it is critical to understand in detail both the local practices of mental health delivery, and the local ideas about mental illness and major services barriers such as stigma.

Structure/Method/Design: Methods of analysis included literature review, policy and service structure analysis, participant observation, and individual as well as focus group interviews with stakeholders from various arenas of the mental health delivery system in Shanghai, China. The methods were formulated within a transdisciplinary framework involving the fields of psychiatry, public health, clinical social work, and medical anthropology in order to gain a comprehensive insight into the Chinese mental health system.

Results (Scientific Abstract)/Collaborative Partners (Programmatic Abstract): A solution narrative of transdisciplinary strategies was generated based on available local resources and strategies acquired from previous field practices. We delineated in detail the service structure, care pathway, and essential skill packages of the mental health delivery system in Shanghai, and the scale of challenges it is facing. The findings generally supported the current Chinese strategy for community mental health service expansion and highlighted multiple barriers to expansion, such as restricted diagnostic privileges, lack of mental health social workers, and stigma experienced in the Chinese culture.

Summary/Conclusion: We concluded that in spite of differences in local context, the experience of Partners in Health in implementing mental health services in low- and middle-income countries would contribute significantly to China's service scale-up efforts. Some initial recommendations include equip community health service centers for the initial diagnosis of common mental disorders; improve task-sharing training for mental health social workers; identify and implement intervention strategies to destigmatize mental illness; and improve community prevention and education programs focused on generating behavioral change. The newly implemented Chinese mental health legislation provides an excellent opportunity for mental health service expansion; however, some aspects of the law should be considered for revision in order to maximize service capacity. Further needs assessment and policy analysis are needed to assess the impact of large-scale internal migration and rapid population aging.

Trauma registries in low- and middle-income countries: Working with what we already have

M.R. Kesinger¹, L. Nagy², A.M. Rubiano³, J.C. Puyana¹, E.W. Etchill⁴; ¹University of Pittsburgh Medical Center, Surgery, Pittsburgh, PA/US, ²University of Pittsburgh, DeKalb, IL/US, ³Universidad Surcolombiana, La Plata, HUILA/CO, ⁴University of Pittsburgh School of Medicine, Department of Surgery, Pittsburgh, PA/US

Background: Trauma registries (TRs) are a fundamental tool of mature trauma systems. They are used in developed countries to measure the effectiveness of trauma quality improvement (TQI). Low- and middle-income countries (LMIC) have not developed TRs on a large scale. Financial barriers and a lack of digital and human

infrastructure are obstacles to creating TRs in LMICs. Some LMIC hospitals have created financial databases to record pay-for-service information. Innovative methods to adapt financial databases into TRs could advance TQIs in LMICs. We report how we adapted a financial database to measure the effectiveness of a TQI.

Structure/Method/Design: A TQI standardizing generally accepted interventions in initial resuscitation of patients at a trauma center in Neiva, Colombia was implemented in September 2011. To measure the effectiveness, we wished to investigate the incidence of standardized interventions in the emergency department (ED) before and after TQI implementation. We used restructuring techniques to create a database that permitted analysis from an administrative hospital database that was used for billing purposes. The database spanned September 2010 to September 2012. To confirm accuracy, a chart review was conducted for a subset of patients by two independent researchers.

Results (Scientific Abstract)/Collaborative Partners (Programmatic Abstract): The restructured database allowed for analysis of mortality, many ED interventions and length of hospital stay 1 year before intervention and 1 year after. Chart review confirmed database fidelity. The analysis demonstrated the general success of the TQI by increased interventions, decreased length of stay, and a dramatic decrease in mortality for severely injured patients. It also demonstrated some interventions that did not increase, showing where future TQI efforts should be directed at the institution. A second TQI is currently underway, based on the results of this work.

Summary/Conclusion: Adapting financial databases into trauma registries is a potentially cheap and effective way to measure trauma quality improvement effectiveness in LMIC hospitals. This method, in turn, could be used as a catalyst to new quality improvement initiatives in LMIC hospitals.

Factors influencing the external validity of the evidence of HIV counseling and testing data in global health settings

E. Marseilli¹, A. Mirzazadeh², M. Malekinejad³, J. Kahn², A.J. Goodell⁴; ¹Health Strategies International, Oakland, CA/US, ²University of California, San Francisco, Institute for Health Policy Studies, San Francisco, CA/US, ³University of California, San Francisco, Epidemiology and Biostatistics, San Francisco, CA/US, ⁴Global Health Sciences, Berkeley, CA/US

Background: Reliable assessments of external validity (EV) are needed in global health decision-making, to ensure that interventions are implemented in settings where favorable benefits observed in evaluation studies can be replicated. To date, there are no standards for how EV is defined or assessed. Ours is the first systematic approach to develop a quantitative tool for assessing EV in global health settings, applied initially to HIV testing and counseling (HTC).

Structure/Method/Design: We conducted a literature search and structured discussions within our team to develop a list of EV indicators (i.e., study characteristics that might determine EV). We grouped indicators into thematic categories. We refined and amended the list through a two-round Delphi process with 28 HTC experts identified as authors of HTC studies. We sent a structured survey to these experts to elicit the weights for each indicator and to propose additional indicators. We sent the first round results back to the experts for re-weighting. We extracted data for EV indicators from 46 HTC studies identified via standard systematic review methods.