

barriers. Barriers may be more complex, relating to: healthcare decision-making, perceptions of quality and traditional culture and beliefs. It is noted that many patients die before reaching care at the facility level.

Going Forward: Multiple methods will provide perspectives from the community on barriers to optimal EC access.

- Case studies: Cases that present to the ED late in the course of illness will be utilized to understand aspects such as recognition of symptoms and severity, first aid administered and deliberations of the patient or caregivers.
- Focus Groups: Emergency scenarios will be utilized to explore the course of action that would be taken by community members.
- Community Survey: Hypothesis will be drawn from the qualitative data and tested with a quantitative survey implemented in locations frequented by a wide cross-section of the community.

This program has potential to increase understanding of rural Ugandan sociocultural factors in EC seeking, impact EC implementation and increase mutual understanding between providers and similar underserved populations.

Funding: This research is not funded.

Abstract #: 2.050_NEP

Assessing diabetes knowledge and prevalence in Nevis, WI: A type 2 diabetes surveillance initiative

J. Richardso¹, Y.P. Tchuisseu¹, D. Omar¹, C. Conn¹, A. Browne², S. King³, M. Crawford-Johnson⁴, N. Swaby², A. Wills⁵, A. Herbert⁴, M. O'Connor⁴, G. Mills²; ¹University of Virginia, Charlottesville, VA, USA, ²Nevis Sixth Form, Nevis, WI, ³Haverford College, Haverford, PA, USA, ⁴Clarence Fitzroy Bryant College, St. Kitts, WI, ⁵City University of New York, New York, NY, USA

Background: A multi-university team investigated the prevalence of type 2 diabetes (DM) and its risk factors in Nevis, WI, the second leading cause of death in the country.

Methods: We used a mixed methodological approach of randomized chart reviews and surveys. The randomized chart review of the six publicly funded health centers yielded 439 examined charts (4% of the island population). Metabolic screenings, medical history and general health information were analyzed using descriptive statistics. The World Health Organization's STEPwise instrument was used to survey 110 Nevis residents (~1% of the island's population) focusing on diabetes knowledge, risk factors (such as diet, exercise, risk-factors), and open-ended questions targeting perceived DM risk.

Findings: Approximately 17% of the clinical records noted a diabetes diagnosis. Distribution of diabetes between males and females was 16% and 17% respectively. Seven percent of 18-49 years olds (n=279) and 34% of 50 – 75 years olds (n=160) are diabetic. Evidence of a family history that included T2D was evident in 62% of records with a family history notation. While not recorded in every file, risk factors of hypertension, elevated cholesterol and fasting blood glucose were evident in 46%, 72% and 51% of the files respectively. Surveys revealed a similar prevalence of diabetes (14%) and persons indicated that clinicians had within the last three years advised them to reduce salt intake (24%), reduce fat in their diet

(34%), reduce their weight (48%) and increase physical activity (43%). Surveys revealed understanding that diet contributed to diabetes (78%) but only 14% indicated inadequate physical activity contributed to DM. While the severity of DM was clear to respondents (e.g. potential amputations), over 1/3 of respondents were not aware of the DM support services offered by the Ministry of Health.

Interpretation: Results suggest that educational materials and interventions targeting DM risk factors would be beneficial if they were widely advertised. Persons in this study also confirmed the MOH's suspicions that type 2 diabetes prevention, early detection and management is a population health concern.

Funding: Financial support for this research was provided by the National Institutes of Health Minority Health International Research training Grant (#5T37MD008659-02).

Abstract #: 2.051_NEP

The burden of orthopaedic disease presenting to a referral hospital in northern Tanzania

Ajay Premkumar, Honest H. Massawe, David J. Mshabaha, Jared R. Foran, Neil P. Sheth

Background: In low and middle-income countries, country specific data is scarce regarding the burden of surgical disease, with most estimates extrapolated from indirect methods. Kilimanjaro Christian Medical Center (KCMC) is the only tertiary referral hospital for a population of over 11 million in Northern Tanzania. This study aims to directly quantify the current orthopaedic burden of disease at KCMC and provide a foundation to estimate the magnitude and potential benefit of improving access to orthopaedic surgical care in the northern regions of Tanzania.

Methods: Prospective data was collected during June 2015 for 113 patients admitted to the Orthopaedic Surgery ward at KCMC. Retrospective review of available hospital records for 11,678 patients presenting to the KCMC Emergency Medicine Department, Orthopaedic Clinic, and Orthopaedic Ward over the previous 12 months was also performed to obtain a more complete picture of the burden of orthopaedic disease seen at this tertiary referral center.

Findings: KCMC treats an average of 11,172 orthopaedic patients each year. Approximately 57.1% of these patients are seen as outpatients in clinic, 30.1% are seen in the emergency department and 12.8% are admitted as inpatients in the orthopaedic ward. Road traffic accidents (RTAs) represented the most common etiology of injury requiring ward admission at 63.7%, followed by falls at 29.2%, and assaults at 4.4%. Of admissions between ages 15-45, 73.5% were from RTAs. The majority of RTAs, 52.8%, involved a motorcycle and 30.56% involved pedestrians. Femur fractures were the most common injury seen (39.0%), followed by tibia (27.2%) and radius (17.7%) fractures. Patients used a wide variety of transportation methods and took 2.3 hours on average to arrive at KCMC from the location of their injury. Once in the hospital, patients averaged a length to surgery of 10.2 days and a hospital length of stay of 13.5 days. Of all admissions, 96.5% had indications for surgical fixation, while only 57.9% received surgery. KCMC is the only tertiary referral center for a five state region; however, 65.7% of patients originated from the same state in which the hospital resides.